

Sc-Cl-H

XII 728 1265

[Se(OH<sub>3</sub>)]<sup>+</sup>(CO<sub>3</sub>)<sup>2-</sup> (Ji, Cerobase over,  
Complexes)

Pactzold R., Garsoffke M.

Z. anorgan. und allgem. Chem., 1915, 836, 51-2,  
52-54 (Recr.; ref. anal).

Untersuchungen an Selen-sulfat-sulf-  
verbindungen. II. Das Selenitacidiumion.  
Raman-Spektren, Kraftwirkungen  
und Bindungsverhältnisse. 10. Erg.-  
Bd. XIX, 1902, 115/160

V. Cl. Cl.<sup>2+</sup>;  $\text{BaCl}_2 \cdot (\text{OH})_2$  (2xK+Y=6) XII 839 1966

Комиссарова Т.И., Москва 5.3.

Февраль. Моск. Ук.-ма, Красн.

1966; № 3, бг-70

10

PX 1964

$\text{SeH}_3\text{Cl}$

Armstrong D.R., uqp..

1972

(γ)

"Theor. Chim. Acta", 1972, 25,  
N 4, 396-9.

(see  $\text{SeH}_3\text{Br}$ , III)

HgSe...HCl

DM. 18547

1984

Hinchliffe fl.,

ab initio  
paerem.      J. Mol. Struct., 1984,  
106, N 3-4, "Theochem";  
15, N 3-4, 361-366.

HCl·H<sub>2</sub>Se (OM 29842) 1988

Macharia N., Sult B. S.,

UK 6  
Mampurie

Inorg. Chem., 1988,  
27, N 13, 2383-2385.

Ch<sub>2</sub>·H<sub>2</sub>Se (0029842) 1988

Macharia N.P., Ault B.S.

UK 6 Inorg. Chem., 1988,  
Maupay 27, (13), 2383-5.

Infrared spectroscopic  
studies of the interaction  
of chlorine fluoride and

chlorine with hydrogen  
selenide, dimethyl selenide  
and arsine in argon  
matrices.