


C<sub>3</sub>F<sub>6</sub>



1963

9551 - P(V<sub>i</sub>)

Brunoy O.V., Miller R.A.,  
 Cousins J.H.

Syntech Molcc. Structure and Spectrosc.  
 Columbus 1963, Columbus, Ohio, Sept  
 59

Vibrational spectra...

 $C_3 \quad F_6$

1965

 $C_3F_6$ 

спектр

19 Б146. Инфракрасные спектры перфторциклоопана и цис- и трансперфторбутенов-2. Heicklen Julian, Wachi Francis, Knight Vester. The infrared spectra of perfluorocyclopropane and cis- and trans-perfluorobutene-2. «J. Phys. Chem.», 1965, 69, № 2, 693—695 (англ.)

Приведены частоты ИК-спектров поглощения в области призмы NaCl трех простых фторуглеводородов — цикло- $C_3F_6$  цис- $C_4F_8$ -2 и транс- $C_4F_8$ -2 и высказаны соображения о соответствии найденных частот определенным колебаниям связей.


Н. Спасокукоцкий

x · 1965

19

☒


1987



Фурсов М. Б.,  
Кошаров В. Н.

Zh. Fiz. Khim.; 41(5), 1065.

Dependence of mass spectra on temperature. II.

Mass spectra of  $C_2F_4$ -(I),  
 $C_2F_3H$ -(II)   $C_3F_6$ -(III).

(See.  $C_2F_4$ ) III

C3-F6

1972

BGP - 4336 - XIV

Lifshitz C. et al.

(2) J. Mass Spectrom. and Ion Phys., 1972, 10, N1, 25-37.

(CIVCF<sub>4</sub>) XIV

60408.7321

Ch,Ex-C,TC,MGU

$C_2F_5^+$  29864  
 $C_2F_6^+$  (3)

8976

# 3-12116

Simm I.G., Danby C.J., Eland J.H.  
D., Mansell P.I.

Translational energy release in the  
loss of fluorine atoms from the  
ions  $SF_6^+$ ,  $CF_4^+$  and  $C_2F_6^+$ .

"J.Chem.Soc.Faraday Trans.", 1976, Part 2,  
72, N 2, 426-434 (англ.)

569 576 585 0593 пик ВИНИТИ

$C_3F_6$  Lommeca 8222 | 1979.

Dowrey A.H., et al.

Moncks r.

Empycnema, J. Mol. Struct., 1979

Ji

53, 189-196.

$C_3 F_6$

1980

Karuppannar T.A., et al

M.N. Indian J. Pure and  
Appl. Phys., 1980, 18, N 8,

● 626 - 629.

(crys.  $C_3 H_6$ ;  $\bar{1}\bar{1}\bar{1}$ )

$\text{CF}_3\text{CF}-\overset{\theta}{\underset{2}{\text{CF}}}$   
neopentylidene  
bis(2-pentenyl)

(Original 12793) 1981.  
Beagley B, et al.

Wacker et.  
Cryogenic

J. Fluorine Chem.,  
1981, 18, 159-171.

$C_3F_6^+$

Lommel 12654

1981.

Berman D. W., et al

Reference

Int. J. Mass Spectrom.

Process., and Ion Phys., 1981

41ff

39, 363-71.

$(CF_3)_2 C$

1982

reuterip.

cmplkmt.

$\Delta E$

Cuthbertson A.F.,  
Glidewell Ch.

of Mol. Struct., 1982,  
87, 11, Theochem., 1982,  
4, 11, 71-79.

(cav.  $(BeH_3)C$ ; III)

:C(CF<sub>3</sub>)<sub>2</sub> (OM 23346) 1986

Dixon D. A.

Железнор. Ж. Phys. Chem., 1986,  
с. 54-56.

90, 54-56.

$F_2C=CF-CF_3$  (OM 28286) 1987


Stavnebrekk P. Y.,  
Corpsen, Bakker P., et al.,  
Meopen  
Patent J. Mol. Struct., 1987, 162  
101-116.

$C_3F_6$

1990

D<sub>0</sub>

113: 221072d Remarks on infrared multiphoton dissociation of perfluoropropene and difluorochloromethane with two wavelengths. Siguenza, C.; Santos, M.; Torresano, J. A.; Pino, A.; Gonzalez-Diaz, P. F. (Inst. Opt., "Daza de Valdes", CSIC, Madrid, Spain 28006). *Spectrochim. Acta, Part A* 1990, 46A(10), 1499-503 (Eng). IR multiphoton dissociation expts. with one or two wavelengths in perfluoropropene and difluorochloromethane were carried out. The addnl. nonresonant wavelength induced an increase of the yield and selectivity of the dissociation process in the larger mol. and an enhancement of the vibrational intermode coupling in the smaller mol. A rather surprising structure in the dissociation yield-pressure plot was discovered for perfluoropropene at the most effective low pressures.



C.A. 1990, 113, N24