

B-N-Bu

V 5018

1965

$(\text{BCl}_2\text{N}_3)_3$, $(\text{BBr}_2\text{N}_3)_3$ (Tm, *ампиртура*)

Paetzold P.I., Gayoso M., Dehnicke K.
Chem. Ber., 1965, 98 (4), 1173-8

Preparation, properties, and vibrational
spectra of the trimeric boron dihaloazides
 $(\text{BCl}_2\text{N}_3)_3$ and $(\text{BBr}_2\text{N}_3)_3$

CA., 1965, 63, N4, 3774g

Be., Ml.

LCM6 JTK F

V-5479

1966


$F_3B_3N_3H_3$, $Cl_3B_3N_3H_3$, $Br_3B_3N_3H_3$
(p, ΔH_v , ΔH_m , ΔS_v , ΔS_m , ΔH_s , ΔS_s)

Laubengayer A.W., Scaife C.W.J.

J.Chem.and Engng Data, 1966, 11, N2, 172-74.

The vapor pressures and some thermodynamic functions of B=terfluoro=, B=trichloro=and B=tri bromoborarine.

RX., 1967, 2/578 Be



$\Delta H_{aq};$

ΔH°_f

~~48~~
-
~~XX~~

Smith B. Ch., et al ¹⁹⁷⁰

Can. J. Chem., 1970,
48, 12, 1936.

(Cell B-C-N) I